


EMBASY **WATERMELON**



**STARKE
AYRES®**



A LARGE TRIPLOID, HYBRID WATERMELON WITH EXCELLENT TASTE

- Ideal for summer and early season
- Large to medium, uniform, seedless fruit
- Deep red, firm flesh with long shelf life
- High brix and excellent taste
- Very high yielding with medium vines

CUSTOMER SERVICES: 0860 782 753 • WWW.STARKEAYRES.CO.ZA • MEMBER OF THE PLENNEGY GROUP

SEEDS OF SUCCESS

*EMBASY

WATERMELON



Type	A hybrid Triploid watermelon.
Maturity	Main season 85 - 95 days.
Plant Characteristics	Vigorous plants with healthy leaf cover.
Fruit Characteristics	Large, uniform, oval fruit with weight of 8 - 11kg. The rind has a medium-dark green colour with light green stripes. The flesh has deep red colour with firm texture, excellent flavour and high brix.
Disease Resistance	Intermediate Resistance: Fusarium wilt (Fon: 1), Co (1).
Climate	Optimum germination temperature: 27°C to 32°C, with night temperature not lower than 24°C. Optimum growth temperatures at night are 18 - 20°C, and day 24 - 30°C, and for ripening. 15 - 25°C. Do not sow or transplant into the open field until the temperatures are warm enough, unless plastic mulches are used.
Soil	Best performance is obtained on newly prepared, sandy-loam soils, which are rich in humus and fer-tilizer, well-drained, with pH (H2O) = 5.6 – 6.8. Heavy soils should be avoided.
Plant Population	2,500 to 3,000 plants/ha, grafted on a suitable rootstock. Interplanted with a suitable pollinator variety (833 to 1,500 plants/ha). Total plant population 3,333 to 4,500 plants/ha.
Fertilisation	Approximate nutrient requirements depending on soil type, soil analyses, and crop load; N: 90-170 kg/ha: 30% pre-plant and the balance as two side dressings. P: 15-180 kg/ha: Apply pre-plant and work down to root depth. K: 40-250 kg/ha: 30% pre-plant and the balance as two side dressings, simultaneously with the N-application (eg. KNO3).
Adaptation	EMBASY is recommended for main season and early production.
Pollination	A suitable pollinator must be inter-planted at a recommended ratio of 3:1 in the main season, when the conditions are favourable. In the early season and periods of difficult pollination, it is advisable to plant at a ratio of 2:1. Consult our seedless production guidelines for further information.
Features and Benefits	EMBASY is very high yielding with uniform fruit size. The flesh has deep red colour, firm texture, excellent flavour and high brix. Suitable for fresh market, fresh-cut and supermarkets.

* Submitted for registration on the national variety list.

INDEMNITY

All technical advice and/or production guidelines given by STARKE AYRES or any of its personnel with reference to the use of its products, is based on the company's best judgement. However, it must be expressly understood that STARKE AYRES does not assume responsibility for any advice given or for the results obtained.

CUSTOMER SERVICES: 0860 782 753 • WWW.STARKEAYRES.CO.ZA • MEMBER OF THE PLENNEGY GROUP

SEEDS OF SUCCESS